

SYNTHESIS AND CHARACTERIZATION OF NOVEL METAL CHELATES OF 2-(8-QUINOLINOL-5-YL) –METHYL AMINO-5-PYRIDINYL-1, 3, 4-THIADIAZOLE DERIVATIVES

Divyesh K Patel^{*1}, Tejas B.Patel², Arun Singh³

^{*1}*Department of Chemistry, Govt. Geetanjali Girls P. G. Collage, Bhopal, M.P, India.*

²*Shree Jayendrapuri Arts and science collage, Bharuch, Gujrat, India*

³*Rajya Shiksha Kendra Pustak Bhawan, Arera Hills, Bhopal, M.P., India*

divyesh.msc@gmail.com

ABSTRACT:

The Synthesis of novel Metal chelates of 5-Chloromethyl-8-quinolinol coupled with 5-pyridinyl-(1,3,4) thiadiazol-2-yl amine has been carried out in the presence of sodium bicarbonate. The newly synthesised compounds were confirmed on the basis of their spectral characterisation like IR, NMR, Mass and their Elemental analysis. The transition metal chelates viz. Cu²⁺, Ni²⁺, Co³⁺, Mn²⁺ and Zn²⁺ of novel ligand were prepared and characterized by metal-ligand (M:L) ratio, IR and reflectance spectroscopic and magnetic properties.

KEYWORDS: Metal chelates, 5-Chloromethyl-8-quinolinol, 5-pyridinyl-(1,3,4) thiadiazol-2-yl amine

INTRODUCTION:

Co-ordination compounds exhibit different characteristic properties which depend on the metal ion to which they are bound. On the basis of nature of the metal as well as the type of ligand these metal complexes have extensive applications in various fields of human interest [I,II]. Chelation or complexation observes more potent antibacterial effect against some microorganisms than the respective drug [III, IV]. Synthesis of Co (II), Ni (II) and Zn(II) complexes with thiazole ring containing Schiff base ligands and their antimicrobial activities were tested against eight different microorganisms [V, VI,VII]. Earlier we have reported biological importance of metal chelates of 5-Chloromethyl-8-quinolinol (CMQ) derivatives coupled with 5-(4-chloroPhenyl)-1, 3, 4-thiadiazol-2-yl amine which reveals that the ligand is moderately toxic against fungi, while all the chelates are more toxic than ligand. Among all the chelates the Cu²⁺ chelate is more toxic against fungi [VIII]. So here in continuation with our earlier work we wish to report synthesis and characterisation of same class of chelates.

EXPERIMENTAL

Synthesis of 2-(8-Quinolinol-5-yl)–methyl amino-5-pyridinyl-1, 3, 4-thiadiazole:

In a round bottom flask, 5-chloromethyl-8-quinolinol hydrochloride (CMQ), (2.3 gm, 0.01 mole) and 2-amino-5-pyridinyl-1, 3, 4-thiadiazole (3.35 gm, 0.01 mole) in acetone (50 ml) were

suspended. To this suspension sodium bicarbonate (1.68 gm, 0.02 mole) was added and the mixture was warmed on the steam bath for about six hours. End of reaction was monitored by TLC. Finally solution made basic with 5% ammonium hydroxide. Final product was collected after recovery of solvent (Acetone). The yellow solid was purified by washing with acetone. Yield of the ligand compound is 77% and having melting point 164°C.

ANALYSIS

C% ,H% ,N% ,S%

Elemental Analysis

Calculated: 60.89 3.88 20.89 9.55

C₁₇H₁₃N₅OS (335)

Found : 60.8 3.7 20.8 9.4

IR Spectral Features: 3400 (NH), 2910 (CH₂), 2850, 1599, 1507, 1450 (aromatic).cm⁻¹)

NMR: ppm 8.51(d 4H pyridinyl), 7.25-7.72 (m 5H Ar-H), 5.76 (OH) 3.35 (CH₂) Signal

Synthesis of metal chelates of novel ligand

The metal chelates of novel ligand with Cu²⁺, Ni²⁺, Co²⁺, Mn²⁺, Zn²⁺ and metal ions were prepared in two steps. All the metal chelates were prepared in an identical procedure. The details are given as follows.

Preparation of compound solution


Ligand (0.05 mol) was taken in 500 mL beaker and formic acid (85% v/v) was added up to slurry formation. To this slurry, water was added till the complete dissolution of compound. It was diluted to 100 mL.

Synthesis of Ligand-metal-chelates

In a solution of metal acetate (0.005 mol) in water (100 mL), 20 ml of above mentioned compound solution (*i.e.* containing 0.01 M compound) was added with vigorous stirring at room temperature. The pH was adjusted around 4.5 to 6 for complete precipitation of metal chelate. The precipitates were digested on a boiling water bath. The precipitates of chelate were filtered off, washed by 1:1 mixture of water: ethanol and finally with acetone and dried at 70°C for 24 hours.

Measurements

The elemental contents were determined by Thermo Finigen Flash1101 EA (Italy), the metals were determined volumetrically by Vogel's method¹⁸. To 100 mg chelate sample, each 1 mL of HCl, H₂SO₄ and HClO₄ were added and then 1 g of NaClO₄ was added. The mixture was evaporated to dryness and the resulting salt was dissolved in double distilled water and diluted to the mark. From this solution the metal content was determined by titration with standard EDTA solution. Infrared spectra of the synthesized compounds were recorded on Nicolet760 FT-IR spectrometer. NMR spectrum of compound was recorded on 400 MHz NMR spectrophotometer. Magnetic susceptibility measurement of the synthesized complexes was carried out on Gouy Balance at room temperature. Mercury tetrathiocyanatocobalate(II) Hg[CO(NCS)₄] was used as a calibrant. The electronic spectra of complexes in solid were recorded on at room temperature. MgO was used as reference.



Scheme 1.

RESULTS AND DISCUSSION

The synthesis of 5-pyridinyl-1,3,4-thiazole-2-yl amino methylene-8-quinolinol was performed by a simple nucleophilic substitution reaction of 5-pyridinyl-1,3,4-thiazole-2-amine and 5-chloromethyl-8-quinolinol hydrochloride (CMQ). The resulted compound was an amorphous yellow powder. The C,H,N,S contents of compound (Table 1) are consistent with the structure predicted (Scheme 1). The IR spectrum of the compound comprises the important bands due to 8-quinolinol. The bands were observed at 1667, 1577, 1508, and 1390 cm⁻¹. The broad band due to -OH group appeared at 3500 cm⁻¹. In this band the inflections are observed at 2989 cm⁻¹. While the latter two might be attributed to asymmetric and symmetric vibration of CH₂ of CMQ. The

NMR spectrum of compound in DMSO indicates that the singlet of 2 H at 3.35 ppm of N-CH₂-Ar group. While the singlet at 5.76 ppm due to -OH group. The aromatic protons are appeared in multiplicity at 7.25 to 7.72 ppm. The pyridinyl protons are appeared in doublet at 8.51 ppm. The vigorous oxidations of compound yield 8-hydroxy quinoline-5-carboxylic acid. The melting point is 164°C. Thus the structure of compound is confirmed as shown in Scheme 1.

The metal and C,H,N contents of metal chelates of compound Table 1 are also consistent with the predicted structure. The results show that the metal: ligand (M:L) ratio for all divalent metal chelate is 1:2.

The infrared spectra of all the chelates are identical and suggest the formation of all the metal cyclic compound by the absence of band characteristic of free -OH group of parent compound. The other bands are almost at their respectable positions as appeared in the spectrum of parent-compound ligand. However, the band due to (M-O) band could not be detected as it may appear below the range of instrument used. The important IR spectral data are shown in Table 2.

Magnetic moments of metal chelates are given in Table 2. The diffuse electronic spectrum of Cu²⁺ chelates shows two broad bands 14970 and 24614 cm⁻¹. The first band may be due to a 2B_{1g} - 1A_{1g} transition, while the second band may be due to charge transfer. The first band shows structures suggesting a distorted octahedral structure for the Cu²⁺ metal chelates. The higher value of the magnetic moment of the Cu²⁺ chelate supports the same. The Co²⁺ metal chelate gives rise to two absorption bands at 24935 and 19897 cm⁻¹, which can be assigned 4T_{1g} - 2T_{2g}, 4T_{1g} - 4T_{1g}(P) transitions, respectively. These absorption bands and the μ_{eff} value indicate an octahedral configuration of the Co²⁺ metal chelate. The spectrum of Mn²⁺ polymeric chelate comprised two bands at 18414 cm⁻¹ and 23782 cm⁻¹. The latter does not have a very long tail. These bands may be assigned to 6 A_{1g} - 4T_{2g}(G) and 6 A_{1g} - 4A_{2g}(G) transitions, respectively. The high intensity of the bands suggests that they may have some charge transfer character.

The magnetic moment is found to be lower than normal range. In the absence of low temperature measurement of magnetic moment it is difficult to attach any significance to this. As the spectrum of the metal chelate of Ni²⁺ show two distinct bands at 22312 and 15716 cm⁻¹ are assigned as 3 A_{2g}(F) - 3T_{1g}(F) and 3 A_{2g}(F) - 3T_{1g}(F) transition, respectively suggested the octahedral environment for Ni²⁺ ion. The observed μ_{eff} values in the range 3.01-3.2 B.M are consistent with the above moiety.

Table 1. Analysis of novel ligand and its metal chelates.

Molecular formula	M.Wt Gm/mole	Yield %	Elemental analysis											
			%Metal		%C		%H		%N		%S			
			Cald.	Found	Cald.	Found	Cald.	Found	Cald.	Found	Cald.	Found		
C ₁₇ H ₁₃ N ₅ O ₅	335	74	--	--	60.89	60.8	3.88	3.7	20.89	20.8	9.55	9.4		
C ₃₄ H ₂₄ N ₁₀ O ₂ S ₂ Cu ⁺² ·2H ₂ O	767.5	74	8.27	8.2	53.15	53.0	3.64	3.5	18.24	18.2	8.33	8.2		
C ₃₄ H ₂₄ N ₁₀ O ₂ S ₂ Ni ⁺² ·2H ₂ O	763	75	7.69	7.6	53.47	53.3	3.66	3.5	18.34	18.2	8.38	8.3		
C ₃₄ H ₂₄ N ₁₀ O ₂ S ₂ Mn ⁺² ·2H ₂ O	759	72	7.23	7.2	53.75	53.6	3.68	3.6	18.44	18.3	8.43	8.4		
C ₃₄ H ₂₄ N ₁₀ O ₂ S ₂ Co ⁺² ·2H ₂ O	763	80	7.72	7.7	53.47	53.4	3.66	3.6	18.34	18.2	8.38	8.3		
C ₃₄ H ₂₄ N ₁₀ O ₂ S ₂ Zn ⁺² ·2H ₂ O	769	77	8.50	8.4	53.05	53.0	3.64	3.6	18.20	18.2	8.32	8.3		

Table 2. Spectral data and magnetic moment of compound metal chelates.

Metal Chelates	μ_{eff}	Electronic spectral data cm^{-1}	Transition	IR spectral features common for all cm^{-1}
Compound-Cu ²⁺	2.14	24614	Charge transfer	1667
		14970	2B1g_2A1g	1577
				1508
Compound-Ni ²⁺	3.35	22312	3A1g_3T1g(P)	1457
		15716	3A1g_3T1g(F)	2889
		24935	4T1g(F)_4T2g(F)	1269
		19897	4T1g(F)_4T2g	3400
Compound-Co ²⁺	5.15	2425	4T1g(F)_4T2g(P)	--
		23782	6A1g_6A2g_4Eg	--
		18414	6A1g_4T2g(4G)	--
		16728	6A1g_4T1g(PG)	--
Compound-Mn ²⁺	5.96			
Compound-Zn ²⁺	--	--	--	--

REFERENCES

- [I] R. Johari, G. Kumar, D. Kumar, S. Singh, Synthesis and Antibacterial Activity of M. (II) Schiff Base Complex, *J. Ind. Coun. Chem*, 2009, 26, 23.
- [II] P. Mittal, V. Uma, Synthesis, spectroscopic and cytotoxic studies of biologically active new Co (II), Ni (II), Cu (II) and Mn (II) complexes of Schiff base hydrazones, *Der Chemica Sinica*, 2010, 1(3). 124.
- [III] H. Kumar, R. Chaudhary, Synthesis and physico-chemical studies of newly formed complexes of transition and inner transition metals with (5-methoxy- 2- {{(4-methoxy-3, 5-dimethy- 1-pyridiny) methyl} sulfinyi}-ih-benzimidazole), *Der Chemica Sinica*, **2010**, 1(2), 55.
- [IV] S.I. Habib, M.A. Baseer, P.A. Kulkarni, Synthesis and physico-chemical studies of newly formed complexes of transition and inner transition metals with (5-methoxy- 2- {{(4-methoxy-3, 5-dimethy- 1-pyridiny) methyl} sulfinyi}-ih-benzimidazole), *Der Chemica Sinica*, 2011, 2(1), 27.
- [V] Sangita Sharma, Jayesh Ramani, Jasmin Bhalodia, Synthesis, Characterization and Antimicrobial Activity of Some Transition Metal Complexes (Mn, Co, Zn, Ni) With L-Proline and Kojic Acid, *Pelagia Research Library*, 2011, 2 (4):374-382
- [VI] Y.J. Thakor, S.G. Patel, K.N. Patel, Synthesis, characterisation and antimicrobial studies on La(III), Ce(III) and Pr(III) complexes with a tetraaza macrocyclic Ligand, *Der Chemica Sinica*, 2011, 2(1), 43.
- [VII] O.W. Salawu, P.K. Onoja, J. F. Sale, Studies on some metal (II) sulphates salts chelates of salicylaldehyde benzoyl acid hydrazone and acetylaldehyde benzoyl acid hydrazone, *Der Chemica Sinica*, 2011, 2(2), 25.
- [VIII] Patel D K, Arun Singh, Synthesis, Characterization and Antimicrobial Activity of Metal Chelates of 5-[4-Chloro phenyl(1, 3, 4) thiadiazol-2-ylamino methylene]-8-hydroxy quinoline, *E-Journal of Chemistry*, 2009, 6(4), 1017-1022

Received on May 4, 2013.