



EXPEDITIOUS GREEN SYNTHESIS OF VERSATILE ORGANIC COMPOUNDS BY DIVERSE METHODS

Bimal Krishna Banik^{*^}

Department of Chemistry and Chemical Biology, Stevens Institute of Technology, Hoboken, New Jersey, USA; The University of Texas M. D. Anderson Cancer Center, 1515 Holcombe Blvd., Houston, Texas 77030, USA; Department of Chemistry, The University of Texas-Pan American, 1201 West University Drive, Edinburg, Texas 78539, USA; ^ Current Address: Community Health Systems of South Texas, Edinburg, Texas 78539, USA; bimalbanik10@gmail.com

Key Words: Green Chemistry, Synthesis, Microwave, Solventless, Water, One-pot, Various reactions

1. Introduction:

Synthesis of organic compounds requires organic solvents and reagents to promote them. Very often, toxic solvents are used for this purpose. Many organic reactions require molar proportion of reactants and reagents to accomplish the goal. Some reactions require longer reaction times and produce products without control of stereochemistry. Therefore, development of facile and effective organic reactions under environmentally friendly conditions is necessary. In this perspective, several reactions are discussed that are performed by microwave-induced reactions, catalytic procedures, in the absence of any solvents, one-pot method and in water. Important reactions are chosen to investigate the feasibility of conducting these reactions under green chemistry conditions.

2. Results:

Microwave-Induced reactions:

The invention of microwave-mediated organic processes in 1986 has opened up significant challenge to diverse scientists and engineers. Microwave oven generates non-ionizing radiation that is responsible for dielectric heating of the reactants [1, 2]. This forces solvents and reactants to cause molecular interactions. At least one component of the reaction must be polar in this process. We have been pursuing microwave-mediated organic reaction for the past 27 years.

Kitchen and programmed microwave oven was used in our endeavor. Many molecules were prepared following microwave-induced reactions [3].

Synthesis of diverse beta lactams was performed by reaction with acid chloride (equivalent) and Schiff base in the presence of N-methylmorpholine or triethylamine. The beta lactams produced had *cis* or *trans* configuration. Diastereoselective and enantioselective reactions were also possible with certain specific substrates. Notably, the yields and stereochemistry of the beta lactams were not the same when microwave-mediated reactions and classical reactions were conducted in some examples. A high power microwave radiation caused to form the more stable product.

Microwave-induced hydrogenolysis of the benzyloxy group in beta lactams to the hydroxy beta lactams group was achieved with ammonium formate and 10% Pd/C. This reaction produced open chain amides due to the cleavage of N-C-4 bond. However, an aromatic ring directly connected to the C-4 position of the beta lactam is necessary for this cleavage reaction to occur. No cleavage of the N-C-4 bond was observed in the absence of an aromatic ring.

Microwave-induced methods have advantages over common heating systems when products are formed through acceleration of reaction following mild conditions, good selectivity and environmentally benign processes. There are some controversies regarding the rate acceleration in the microwave oven. Some authors believe it is the because of the rapid rise of temperature a reaction rate is enhanced. Some authors believe non-thermal effects are responsible for the acceleration of the reaction rate in most of the examples.

Stereoselectivity of beta lactam formation:

Stereochemistry plays the central role in controlling the medicinal activities of organic compounds. Development of stereospecific reactions, therefore, is the major challenge. Beta lactams are powerful medicines and medicinal activities depend on relative and absolute stereochemistry. Stereochemistry control in beta lactam synthesis is a part of green chemistry process. The formation of a single stereoisomer eliminates purification by chromatography and thus excessive solvents requirement for separation. It was found that *cis* hydroxy beta lactam derivatives are obtained at 0-5⁰C by cycloaddition of acid chlorides and imines in the presence of a tertiary base (triethylamine). The same reaction at high temperature produced *trans* hydroxy beta lactam derivatives in major proportion (over 90%). In addition, by changing the conditions of the experiments, it was also possible to control stereospecificity of beta lactam synthesis. For example, a slow addition of a tertiary base (triethylamine or N-methylmorpholine) to a refluxing solution of a diarylimine with acid chlorides in toluene produced *trans* beta lactam. On the other hand, imines derived from carbohydrates and primary amines produced optically active *cis* beta lactams exclusively regardless of the choice of solvent. The stereochemical results remained identical when microwave irradiation was used for the preparation of these beta lactam derivatives. It was seen that low power microwave irradiation produces *cis* beta lactam in higher proportions than *trans* beta lactams. However, high power microwave-induced reactions of the same substrates produced *trans* beta lactams as major proportion. Pure *cis* beta lactams on exposure to high power microwave irradiation in the presence of tertiary base did not isomerizes

to *trans* beta lactams indicating that that the pathways for the formation of thermodynamically more stable *trans* beta lactams is different than *cis* beta lactam formation [3a, d, g, j, k].

Synthesis of Heterocycles by Microwave-induced Reactions:

Microwave-induced iodine-catalyzed reaction was performed for the synthesis quinazolines via condensation of 1, 2-dicarbonyl compounds and 1, 2-diamines [4a]. Tetrahydroquinolines were prepared *via* microwave induced reactions [4b]. Molecular rearrangement of beta-lactams to pyrrolidine was performed in methanol using microwave-induced reaction [4c]. Microwave-Assisted preparation of pyridinyl-substituted quinoline was achieved through bismuth nitrate-catalyzed Diels-Alder reaction [4d]. Bismuth nitrate-catalyzed microwave-mediated route for the synthesis of dihydropyridines was realized [4e]. Octahydroxanthenes were prepared through bismuth iodide-catalyzed microwave-induced reaction [4f]. Ammonium chloride was found to be an excellent choice for the preparation of diverse pyrroles under microwave-induced conditions [4g, 4h].

Synthesis of Indoles by Microwave-induced Reactions:

Bismuth nitrate was found to be an excellent catalyst for the electrophilic substitution reaction of 7-aza-indole with activated carbonyl compounds under microwave-mediated conditions [5a]. This reaction was slow because of the less reactivity of the 7-aza-indole system. In contrast, the same reaction with indoles proceeded fast [5b]. Microwave-induced ruthenium chloride-catalyzed reaction of isatin with 4-hydroxy proline derivatives produced 3-pyrrole substituted indoles in excellent yield [5c].

Solventless Reactions:

Reactions described above were also performed in the absence of solvents. It was gratifying to note that these reactions are completed within a few minutes instead of hours that are required in the presence of solvents. Iodine-catalyzed Michael reactions of indoles with unsaturated carbonyl compounds were performed under solventless conditions [6a]. Bismuth nitrate-catalyzed synthesis of several *bis* indole derivatives were prepared in excellent yields [6b]. A facile synthesis of N-substituted pyrrole derivatives were conducted *via* bismuth nitrate-catalyzed reaction in the absence of any solvent [6c]. These pyrrole syntheses were also performed by iodine-catalyzed reaction in the absence of solvent [6d]. Polyhydroquinoline was prepared in the absence of solvent *via* microwave-induced reaction [6e]. Pechmann reaction was performed in the absence of any solvent *via* bismuth nitrate-catalyzed microwave-mediated reaction [6f]. Bismuth nitrate-catalyzed microwave-induced simple enamination of β -dicarbonyl compounds was performed in the absence of solvent [6g].

Reactions in Water:

Many organic reactions were performed in water. For example, bismuth nitrate-catalyzed Michael reaction of indoles with unsaturated ketones was conducted successfully in water [7a]. Aza-Michael reaction was also done in water [7b]. Dowex-50 was found to catalyze synthesis of dihydropyrimidines in water [7c]. Indium metal-catalyzed Strecker reaction of imines was

conducted in water successfully [7d]. Phosphoric acid-catalyzed Aza-Michael reaction was performed in water [7e]. This reaction was also conducted in water in the presence of ultrasound [7f]. Diverse pyrroles were prepared following Paal-Knorr method in aqueous carbon dioxide [7g, 7h].

One-Pot method:

Synthesis of new dibenzofluorene derivatives was performed in a one-pot operation. Thus, reaction of 2-naphthylbromide with beta tetralone in the presence of sodium hydride afforded the alkylated ketone. This ketone on cyclodehydration and aromatization reaction in a one-pot method produced dibenzofluorene in good yield. An extension with methoxy aromatic bromide was also possible for the preparation of penatcyclic derivative. These hydrocarbons were converted to numerous compounds that have demonstrated promising anticancer activities [8]

Various Reactions:

Our group had also investigated numerous reactions under green chemistry conditions. For example, bismuth nitrate-mediated microwave-induced selective hydrolysis of amide was performed [9a]. Diels Alder reaction was investigated under an identical condition with great success [9b]. N-bromosuccinimide and polystyrenesulfonate-catalyzed microwave-assisted synthesis of various pyrroles were achieved [9c, 9d, 9e]. Microwave-assisted bismuth nitrate-induced oxidation of curcumin to vanillin was performed [9f]. Aza-Diels Alder reaction proceeded in the presence of bismuth nitrate and microwave irradiation [9g].

Acknowledgment:

The author is grateful to NIH, NCI, Kleberg Foundation, Stevens Institute of Technology, University of Texas M. D. Anderson Cancer Center, University of Texas Health Science Center at San Antonio and University of Texas-Pan American.

References:

1.(a) Gedye, R.; Smith, F.; Westaway, K.; Ali, H.; Baldisera, L; Laberge, L.; Rousell, J.; *Tetrahedron Letters*; **1986**, 27, 279-282; (b) Bose, A. K.; Manhas, M. S.; Ghosh, M.; Shah, M.; Raju, V. S.; Bari, S. S.; Newaz, S. N.; Banik, B. K.; Barakat, K. J.; Chaudhury, A. G.; "Microwave-Induced Organic Reaction Enhancement Chemistry. 2. Simplified Techniques"; *J. Org. Chem.*, **1991**, 56, 6968-6970.

2.Loupy, A. (ed), Wiley-VCH, Weinheim, 2006.

3. (a) Banik, B. K.; Manhas, M. S.; Kaluza, Z.; Barakat, K. J.; Bose, A. K.; "Microwave-Induced Organic Reaction Enhancement Chemistry. 4 Convenient Synthesis of Enantiopure α -Hydroxy- β -lactams"; *Tetrahedron Lett.*, **1992**, 33, 3603-3606; (c) Bose, A. K.; Banik, B. K.; Barakat, K. J.; Manhas, M. S.; "Simplified Rapid Hydrogenation Under Microwave Irradiation: Selective Transformations of β -Lactams"; *Synlett*; **1993**, 575-577; (d) Banik, B. K.; Manhas, M. S.; Newaz, S. N.; Bose, A. K.; "Facile Preparation of Carbapenem Synthons via Microwave-

Induced Rapid Reaction”; *Bioorg. & Med. Chem. Lett.*, **1993**, 3, 2363-2368; (e) Bose, A. K.; Manhas, M. S.; Banik, B. K.; Robb, E. W.; “Microwave-Induced Organic Reaction Enhancement (MORE) Chemistry: Techniques for Rapid, Safe and Inexpensive Synthesis”; *Res. Chem. Interm.*, **1994**, 20, 1-11; (f) Banik, B. K.; Barakat, K. J.; Wagle, D. R.; Manhas, M. S.; Bose, A. K.; “Microwave-Assisted Rapid and Simplified Hydrogenation”, *J. Org. Chem.*, **1999**, 64, 5746-5753; (g) Banik, B. K.; Aguilar, H.; Cordova, D.; “Unprecedented Stereocontrol of β -Lactam Formation Derived From N-Cinnamylidene Arylamine”, *Heterocycles*; **2008**, 11, 2321-2329; (i) Ramos, K.; Banik, B. K.; “Microwave-Induced Clay-Mediated Preparation of Imines: One-Pot Synthesis of β -Lactams”, *Heterocyclic Letters*; **2011**, 27-30; (j) Banik, I.; Becker, F. F.; Banik, B. K.; “Microwave-Induced Stereospecific Synthesis of β -Lactams Derived from Polyaromatic Imines: Influence of Multicyclic Rings at the Nitrogen”, *Heterocyclic Letters*; **2011**, 55-57; (k) Bandyopadhyay, D.; Yanez, M. A.; Banik, B. K.; “Microwave-Induced Stereoselectivity of β -Lactam Formation, Effects of Solvents”; *Heterocyclic Letters*; **2011**, 65-67; (n) Bandyopadhyay, D.; Cruz, J.; Banik, B. K.; “Microwave-Induced Synthesis of 3-Pyrrole Substituted β -Lactams Via Bismuth Nitrate-Catalyzed Reaction”, *Tetrahedron Symposium-in-Print*; **2012**, 68, 10686-10695. Also see: (a) Bose, A. K.; Banik, B. K.; Newaz, S. N.; Manhas, M. S.; “Vinyl β -Lactams: Convenient Elaboration of The Thienamycin Side Chain”, *Synlett*, **1993**, 897-899; (b) Banik, B. K.; Manhas, M. S.; Newaz, S. N.; Bose, A. K.; “Facile Preparation of Carbapenem Synthons Via Microwave-Induced Rapid Reaction”, *Bioorg. & Med. Chem. Lett.*, **1993**, 3, 2363-2368; (c) Manhas, M. S.; Banik, B. K.; Mathur, A.; Vincent, J.; Bose, A. K.; “Microwave-Assisted Synthesis of Vinyl β -Lactam: Synthons for Natural Products”, *Tetrahedron* **2000**, 56, 5587-5601; (d) Bandyopadhyay, D.; Banik, B. K. “Microwave-Induced Stereoselectivity of β -Lactam Formation with Dihydrophenanthrenyl Imines via Staudinger Reaction”, *Helv. Chim. Acta.*, **2010**, 93, 298-301; (e) Banik, I.; Samajdar, S.; Banik, B. K. “Microwave-Induced Stereospecific Synthesis of β -Lactams Derived from Polyaromatic Imines: Influence of Multicyclic Rings at the Nitrogen”, *Heterocyclic Letters*, **2011**, 55-57; (f) Bandyopadhyay, D.; Cruz, J.; Banik, B. K. “Microwave-Induced Synthesis of 3-Pyrrole Substituted β -Lactams Via Bismuth Nitrate-Catalyzed Reactions”, *Tetrahedron Symposium-in-Print*, **2012**, 68, 10686-10695; (g) Bandyopadhyay, D.; Yanez, M.; Banik, B. K. “Microwave-Induced Stereoselectivity of β -Lactam Formation: Effects of Solvents”, *Heterocyclic Letters*, **2011**, 65-67; (h) Ramos, K.; Banik, B. K.; “Microwave-Induced Clay-Mediated Preparation of Imines: One-Pot Synthesis of β -Lactams”, *Heterocyclic Letters*, **2011**, 27-30; (i) Banik, I.; Yadav, R. N.; Banik, B. K. “Microwave-Induced Synthesis of Bis β -Lactams from Hydrazides”, **2017**, *Asian J. Org. & Med. Chem.*, in press; (j) Banik, I.; Becker, F. F.; Banik, B. K.; “Microwave-Mediated Chiral Synthesis of O-Glycosides of β -Lactams”, *Asian J. Org. & Med. Chem.*, **2017**, in press; (k) Banik, I.; Samajdar, S.; Becker F. F.; Banik, B. K. “Microwave-Induced Synthesis of O-Glycosides: Precursors for Anticancer β -Lactams”, *Asian J. Org. & Med. Chem.*, **2017**, in press.

4. (a) Bandyopadhyay, D.; Mukherjee, S.; Rodriguez, R.; Banik, B. K.; “An Effective Microwave-Induced Iodine-Catalyzed Method for the Synthesis of Quinoxalines via Condensation of 1,2-Diamines with 1,2-Dicarbonyl Compounds”, *Molecules*, **2010**, 15, 4207-4212; (b) Banik, B. K.; “Sterospecific Synthesis of Tetrahydroisoquinolines via Microwave-Induced Reaction”, *Heterocyclic Letters*, **2011**, 49-51; (c) Banik, I.; Samajdar, S.; Banik, B. K.; “Microwave-Induced Sodium-Methoxide-Mediated Molecular Rearrangements of β -Lactams to 3-Substituted Pyrrolidines”, *Heterocyclic Letters*, **2011**, 69-72; (d) Rodriguez, R.; Gomez, K.;

Banik, B. K.; “Microwave-Assisted Synthesis of Pyridinyl Substituted Quinoline Through Bismuth Nitrate-Catalyzed Diels-Alder Reaction”, *Heterocyclic Letters*, **2011**, 93-94; (e) Bandyopadhyay, D.; Maldonado, S.; Banik, B. K. “Microwave-Assisted Bismuth Nitrate-Catalyzed Unique Route Toward Dihydropyridines”, *Molecules*, **2012**, 17, 2643-2662; (f) Chavez, A.; Cruz, J.; Munoz, A.; Yadav, R. N.; Bandyopadhyay, D.; Banik, B. K.; “Microwave-Induced Bismuth Iodide-Catalyzed Synthesis of Octahydroxanthenes”, *Heterocyclic Letters*, **2017**, 7, 507-511; (g) Bandyopadhyay, D.; Banik, B. K.; “Microwave-Induced Paal-Knorr Reaction With Ammonium Chloride: Synthesis of Pyrroles”, *Heterocyclic Letters*, **2017**, 7, 473-474; (h) Bandyopadhyay, D.; Banik, B. K.; “Ammonium Chloride-Induced Synthesis of Pyrroles Via Paal-Knorr Reaction”, *Heterocyclic Letters*, **2017**, 7, 463-465.

5. (a) Rivera, S.; Bandyopadhyay, D.; Banik, B. K.; “Microwave-Induced Bismuth Nitrate-Catalyzed Electrophilic Substitution of 7-Aza Indole with Activated Carbonyl Compounds Under Solvent-Free Conditions”, *Heterocyclic Letters*, **2011**, 43-46; (b) Rivera, S.; Iglesias, L; Bandyopadhyay, D.; Banik, B. K.; “Microwave-Induced Bismuth Nitrate-Catalyzed Electrophilic Substitution of Indole with Keto Ester Under Solvent-Free Conditions”, *Heterocyclic Letters*, **2011**, 73-74; (c) Bandyopadhyay, D.; Banik, A.; Batta, S.; Banik, B. K.; “Microwave-Assisted Ruthenium Trichloride-Catalyzed Synthesis of Pyrrole Fused with Indolinone”, *Heterocyclic Communications*, 2009, 121-122R.

6. (a) Banik, B. K.; Farnandez, M.; Alvarez, C.; “Iodine-Catalyzed Michael Reaction of Indoles Under Solventless Conditions”, *Tetrahedron Lett.*, **2005**, 46, 2479-2482; (b) Iglesias, L; Aguilar, C.; Bandyopadhyay, D.; Banik, B. K.; “Bismuth Nitrate-Catalyzed Synthesis of Bis Indolylmethane Under Solventless Conditions”, *Synthetic Communications*, **2010**, 40, 3678-3682; (c) Rivera, S.; Banyopadhyay, D.; Banik, B. K.; “Facile Synthesis of N-Substituted Pyrroles Via Microwave-Induced Bismuth Nitrate-catalyzed Reaction Under Solventless Conditions”, *Tetrahedron Lett.*, **2009**, 50, 5445-5448; (d) Bandyopadhyay, D.; Mukherjee, S.; Banik, B. K.; An Expeditious Synthesis of N-Substituted Pyrroles by Microwave-Induced Iodine-Catalyzed Reactions Under Solventless Conditions; *Molecules*; **2010**, 15, 2520-2525; (e) Castillo, M.; Ortiz, M.; Banik, B. K.; “Microwave-Assisted Synthesis of Polyhydroquinoline in the Absence of Solvent”, *Heterocyclic Letters*, **2011**, 89-91; (f) Aguilar, H.; Reddy, A.; Banik, B. K.; “Microwave-Induced Bismuth Nitrate-Catalyzed Pechman Reaction Under Solventless Condition”, *Heterocyclic Letters*, **2011**, 95-96; (g) Bandyopadhyay, D.; Banik, B. K.; “Microwave-Induced Bismuth Nitrate-Catalyzed Expeditious Enamination of β -Dicarbonyl Compounds Under Solvent-Free Conditions”, *Heterocyclic Letters*, **2011**, 17-21.

7. (a) Banik, B. K.; Garcia, I.; Morales, F.; “Bismuth Nitrate-Catalyzed Michael Reaction of Indoles in Water”, *Heterocycles*, **2007**, 71, 919-924; (b) Kall, A.; Bandyopadhyay, D.; Banik, B. K.; “Microwave-induced Aza-Michael Reaction in Water: A Remarkable Simple Procedure”, *Synthetic Communications*, **2010**, 40, 1730-1735; (c) Mukhopadhyay, C.; Datta, A.; Banik, B. K.; “Dowex-50 in Aqueous Medium: Highly Efficient Biginelli Condensation for the Synthesis of 4-Aryl-3,4-dihydropyrimidones”, *Heterocycles*, **2007**, 71, 181-188; (d) Bandyopadhyay, D.; Velaquez, J.; Banik, B. K.; “Indium-Catalyzed Strecker Reaction in Water”, *Org. & Med. Chem. Lett.*, **2011**, 1, 11-15; (e) Bandyopadhyay, D.; Maldonado, S.; Banik, B. K.; “Phosphoric Acid-Catalyzed Aza Michael Reaction in Water”, *Heterocyclic Letters*, **2011**, 13-16; (f) Bandyopadhyay, D.; Mukherjee, S.; Turrubiarres, L.; Banik, B. K.; “Ultrasound-Assisted Aza-

Michael Reaction in Water”, *Ultrasonics Sonochemistry*, 2012, 19, 969-973; (g) Yadav, R. N.; Banik, B. K.; “Carbon Dioxide-Mediated Preparation of Pyrroles in Water Following Paal-Knorr Method”, *Heterocyclic Letters*, **2017**, 7, 905-906. (h) Yadav, R. N.; Banik, B. K. “Novel Synthesis of Pyrroles via Carbon Dioxide-Mediated Reaction”, *Heterocyclic Letters*, **2017**, 7, 893-894.

8. (a) Banik, B. K.; Becker, F. F.; “Synthesis, Electrophilic Substitution and Structure-Activity Relationship Studies of Polycyclic Aromatic Compounds for the Development of Anticancer Agents”, *Curr. Med. Chem.* **2001**, 8, 1513-1533; (b) Becker, F. F.; Mukhopadhyay, C.; Hackfeld, L.; Banik, I.; Banik, B. K.; “Polycyclic Aromatic Compounds as Anticancer Agents: Synthesis and Biological Evaluation of Dibenzofluorene Derivatives”, *Bioorg. Med. Chem.*, **2000**, 8, 2693-2699; (c) Banik, B. K.; Mukhopadhyay, C.; Becker, F. F. “An Expedient One-Pot Synthesis of Hydrofluorene Derivatives”, *Synthetic Communications*, **2001**, 31, 2399-2403; (d) Banik, B. K. “A Personal Perspective on Medicinal and Pharmaceutical Chemistry”, *Frontiers in Medicinal and Pharmaceutical Chemistry*, doi:10.3389/fchem. **2014.00008**; (e) Becker, F. F.; Banik, B. K., “Synthesis and Biological Evaluation of Novel Dibenzofluorenes”, *Frontiers in Medicinal and Pharmaceutical Chemistry*, 2:55; DOI: 10.3389/fchem.2014.00055; **2014**.

9. (a) Bandyopadhyay, D.; Fonseca, R. S.; Banik, B. K.; “Microwave-Induced Bismuth Nitrate-Mediated Selective Hydrolysis of Amide”, *Heterocyclic Letters*, **2011**, 75-77; (b) Mukherjee, S.; Solano Fonseca, R.; Danso, R.; Banik, B. K.; “Microwave-Induced Bismuth Nitrate-Catalyzed Intramolecular Diels-Alder Reaction”, *Heterocyclic Letters*, **2011**, 87-88; (c) Andoh-Baidoo, R.; Mukherjee, S.; Banik, B. K.; “Microwave-Induced N-Bromosuccinimide-Mediated Novel Synthesis of Pyrroles”, *Heterocyclic Letters*, **2011**, 103-106; (d) Vargas, R.; Leal, B.; Reddy, A.; Bandyopadhyay, D.; Banik, B. K.; “Microwave-Assisted Polystyrenesulfonate-Catalyzed Synthesis of Novel Pyrroles”, *Org. & Med. Chem. Lett.*, **2012**, 2: 24-30 (doi:10.1186/2191-2858-2-24); (e) Andoh-Baidoo, R.; Danso, R.; Mukherjee, S.; Bandyopadhyay, D.; Banik, B. K.; “Microwave-Induced N-Bromosuccinimide-Mediated Novel Synthesis of Pyrroles Via Paal-Knorr Reaction”, *Heterocyclic Letters*, **2011**, 107-109; (f) Bandyopadhyay, D.; Banik, B. K.; “Bismuth Nitrate-Induced Microwave-Assisted Expedient Synthesis of Vanillin From Curcumin”, *Org. & Med. Chem. Lett.*, **2012**, 2, 15-18 (doi:10.1186/2191-2858-2-15); (g) Yadav, R. N.; Reddy, A.; Banik, B. K.; “Bismuth Nitrate-Catalyzed Microwave-Assisted Aza Diels-Alder Reaction for the Synthesis of Bicyclo (2,2,2)-octanones Scaffold”, *Current Microwave Chemistry*, **2014**, 1, 94-97.